

# NLO contributions in the pQCD approach

Zhen-Jun Xiao

*The Department of Physics and Institute of Theoretical Physics, Nanjing Normal University, Nanjing, Jiangsu 210097, P.R. China*

The NLO contributions to some two-body charmless hadronic B meson decays, such as those from the QCD vertex corrections, the quark loops and the chromo-magnetic penguins, have been calculated in the pQCD factorization approach.

## 1. INTRODUCTION

The two-body charmless hadronic B meson decays play an important role in testing the standard model (SM) and probing for the new physics beyond the SM. At present, the key point is how to reduce still large theoretical uncertainty in evaluating the hadronic matrix elements  $\langle M_2 M_3 | O_i | B \rangle$ .

During the past decade, many  $B \rightarrow M_2 M_3$  decays have been studied by employing the QCD factorization approach [1], the soft collinear effective theory (SCET) [2] and the perturbative QCD (pQCD) factorization approach [3, 4]. In the pQCD approach, some next-to-leading order (NLO) corrections, such as those from vertex QCD corrections, the quark-loops and the chromo-magnetic penguins, have been included recently [5, 6, 7], but much more NLO contributions should be calculated in order to give more reliable pQCD predictions.

## 2. OUTLINE OF THE PQCD FACTORIZATION APPROACH

In the pQCD approach, the decay amplitude  $\mathcal{A}(B \rightarrow M_2 M_3)$  can be written conceptually as the convolution,

$$\mathcal{A} \sim \int d^4 k_1 d^4 k_2 d^4 k_3 \text{Tr} [C(t) \Phi_B(k_1) \Phi_{M_2}(k_2) \Phi_{M_3}(k_3) H(k_1, k_2, k_3, t)], \quad (1)$$

where  $k_i$ 's are momenta of light quarks included in each meson, and Tr denotes the trace over Dirac and color indices. The Wilson coefficient  $C(t)$  is evaluated at hard scale  $t$ . The function  $H(k_1, k_2, k_3, t)$  describes the hard dynamics and could be calculated perturbatively. The wave function  $\Phi_{M_i}$  describes hadronization of the quark and anti-quark in the meson  $M_i$ . Using the light-cone coordinates the B meson momentum  $P_B$  and the two final state meson's momenta  $P_2$  and  $P_3$  (for  $M_2$  and  $M_3$  respectively) can be written as

$$P_B = \frac{M_B}{\sqrt{2}}(1, 1, \mathbf{0}_T), \quad P_2 = \frac{M_B}{\sqrt{2}}(1 - r_3^2, r_2^2, \mathbf{0}_T), \quad P_3 = \frac{M_B}{\sqrt{2}}(r_3^2, 1 - r_2^2, \mathbf{0}_T), \quad (2)$$

where  $r_i = m_i/M_B$ .  $m_2$  and  $m_3$  are the mass of the two final state mesons. Putting the anti-quark momenta in B,  $M_2$  and  $M_3$  meson as  $k_1$ ,  $k_2$ , and  $k_3$ , respectively, we can choose

$$k_1 = (x_1 P_1^+, 0, \mathbf{k}_{1T}), \quad k_2 = (x_2 P_2^+, 0, \mathbf{k}_{2T}), \quad k_3 = (0, x_3 P_3^-, \mathbf{k}_{3T}). \quad (3)$$

Then, the integration over  $k_1^-$ ,  $k_2^-$ , and  $k_3^+$  in eq.(1) will lead to

$$\mathcal{A} \sim \int dx_1 dx_2 dx_3 b_1 db_1 b_2 db_2 b_3 db_3 \cdot \text{Tr} [C(t) \Phi_B(x_1, b_1) \Phi_{M_2}(x_2, b_2) \Phi_{M_3}(x_3, b_3) H(x_i, b_i, t) S_t(x_i) e^{-S(t)}], \quad (4)$$

where  $b_i$  is the conjugate space coordinate of  $k_{iT}$ ,  $S_t(x_i)$  is the threshold function, and  $e^{-S(t)}$  is the Sudakov form factor which suppresses the soft dynamics effectively.




 Figure 1: The typical Feynman diagrams which may contribute to the  $B \rightarrow M_2 M_3$  decays at leading order.


Figure 2: The NLO contributions in pQCD approach from the vertex QCD corrections (a-d), the quark-loops (e-f) and the chromo-magnetic penguins (g-h).

### 3. LO and NLO CONTRIBUTIONS

At the leading order in pQCD approach, there are three type diagrams contributing to the  $B \rightarrow M_2 M_3$  decays, the factorizable emission diagrams, the hard-spectator diagrams and the annihilation diagrams, as illustrated in Fig.1. From the factorizable emission diagrams Fig.1a-1d, the corresponding form factors can be extracted by perturbative calculation. For a given decay mode, one can find firstly the individual decay amplitude by evaluating analytically the corresponding Feynman diagram, obtain the total decay amplitude by summing up the different pieces and finally calculate the branching ratios and the CP violating asymmetries.

When compared with the previous LO calculations in pQCD, the following NLO contributions should be considered:

1. The NLO Wilson coefficients  $C_i(M_W)$ , and the NLO RG evolution matrix  $U(t, m, \alpha)$  as defined in Ref. [8], will be used.
2. The strong coupling constant  $\alpha_s(t)$  at two-loop level will be used.
3. Besides the LO hard kernel  $H^{(0)}(\alpha_s)$ , the NLO hard kernel  $H^{(1)}(\alpha_s^2)$  should be included. All the Feynman diagrams, which lead to the decay amplitudes proportional to  $\alpha_s^2(t)$ , should be considered. Such Feynman diagrams can be grouped into the following classes:

I: The NLO contributions from the vertex QCD corrections, the quark-loops and chromo-magnetic penguins, as illustrated in Fig. 2. For chromo-magnetic penguins, there are totally nine relevant Feynman




Figure 3: The four typical Feynman diagrams, which contributes to the form factors at NLO level.




Figure 4: The four typical hard-spectator Feynman diagrams, which contributes at NLO level.




Figure 5: The four typical annihilation Feynman diagrams, which contributes at NLO level.

diagrams, but the first two ( Fig.2g-2h ) provide the dominant NLO contributions [9].

II: The NLO contributions to the LO Feynman diagrams (1a,1b) will affect the extraction of the form factors, as illustrated in Fig. 3. There are totally 13 relevant Feynman diagrams, we here show four of them only.

III: The NLO contributions to the LO hard-spectator Feynman diagrams (1c,1d), as illustrated in Fig. 4. There are totally 56 relevant Feynman diagrams, we here show four of them only.

IV: The NLO contributions to the annihilation Feynman diagrams (1e,1h), as illustrated in Fig. 5. we here show only four of them.

At present, the calculations for the vertex corrections, the quark-loops and chromo-magnetic penguins have been available [5, 6, 7]. For more details about the pQCD predictions for the branching ratios and CP violating asymmetries of  $B \rightarrow K\eta^{(\prime)}, KK^*, \rho(\omega, \phi)\eta^{(\prime)}$  decays, one can see Refs. [6, 7]. It is an urgent task to calculate the Feynman diagrams as shown in Figs. 3-5, in order to provide pQCD predictions at full NLO order.

## Acknowledgments

Work supported by the National Natural Science Foundation of China under Grant No.10575052 and 10735080.

## References

- [1] M. Beneke, G. Buchalla, M. Neubert, and C.T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999); M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
- [2] C.W. Bauer et al., Phys. Rev. D **70**, 054015 (2004); M. Beneke and D. Yang, Nucl. Phys. B **736**, 34 (2006).
- [3] H.N. Li, Prog. Part. & Nucl. Phys. **51**, 85 (2003) and references therein.
- [4] X. Liu, H.S. Wang, Z.J. Xiao, L.B. Guo, and C.D. Lü, Phys. Rev. D **73**, 074002 (2006); A. Ali et al., Phys. Rev. D **76**, 014008 (2007).
- [5] H.N. Li, S. Mishima, A.I. Sanda, Phys. Rev. D **72**, 114005 (2005); H.N. Li and S. Mishima, Phys. Rev. D **74**, 094020 (2006).
- [6] Z.Q. Zhang and Z.J. Xiao, arXiv:0807.2022[hep-ph]; and 0807.2024[hep-ph].
- [7] Z.J. Xiao, Z.Q. Zhang, X. Liu and L.B. Guo, Phys. Rev. D **78**, 114001 (2008); arXiv: 0807.4265[hep-ph].
- [8] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev.Mod.Phys. **68**, 1125 (1996).
- [9] S. Mishima and A.I. Sanda, Prog. Theor. Phys. **110**, 549 (2003).